



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The general health of Barbados continues good, and even though the hot season is well advanced there is no increase in the general sickness. The mosquitoes have materially increased during the past month and many of the *Stegomyia fasciata* and of the *Culex fatigans* may be found. There are no diseases of a quarantinable nature now present on this island.

Report from Castries, St. Lucia Island—Transactions of Service—Mortality.

Passed Assistant Surgeon Fricks reports, July 1, as follows:

Week ended July 1, 1905:

Number of bills of health issued: Supplemental 5, original 1.

Number of vessels fumigated, none.

The sanitary condition of the port continues good.

There were during the month of June 44 deaths, none of them due to a quarantinable disease.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

AUSTRALIA—*New South Wales—New Castle.*—Month of May, 1905. Estimated population, 50,600. Total number of deaths, 51, including diphtheria 1, plague 2, and 5 from tuberculosis.

BAHAMAS—*Dunmore Town.*—Two weeks ended June 26, 1905. Estimated population, 1,232. One death. No contagious diseases.

Governor's Harbor.—Week ended July 1, 1905. Estimated population, 1,500. No deaths and no contagious diseases.

Green Turtle Cay—Abaco.—Two weeks ended June 29, 1905. Estimated population, 3,314. No deaths and no contagious diseases reported.

Nassau.—Two weeks ended July 1, 1905. Estimated population, 12,650. Number of deaths not reported. No contagious diseases reported.

CANADA—*Quebec, Sherbrooke.*—Month of June, 1905. Estimated population, 13,000. Total number of deaths, 22, including 1 from whooping cough.

GREAT BRITAIN—*England and Wales.*—The deaths registered in 76 great towns in England and Wales during the week ended June 24, 1905, correspond to an annual rate of 13 per 1,000 of the aggregate population, which is estimated at 15,609,377.

London.—One thousand and forty-seven deaths were registered during the week, including measles 38, scarlet fever 7, diphtheria 11, whooping cough 26, enteric fever 3, and 14 from diarrhea. The deaths from all causes correspond to an annual rate of 11.7 per 1,000. In Greater London 1,475 deaths were registered. In the "outer ring" the deaths included 4 from measles, 3 from scarlet fever, 5 from whooping cough, and 3 from diarrhea.

Ireland.—The average annual death rate represented by the deaths registered during the week ended June 24, 1905, in the 21 principal town districts of Ireland was 16 per 1,000 of the population, which is estimated at 1,093,959. The lowest rate was recorded in Wexford, viz, 4.7, and the highest in Clonmel, viz, 35.9, per 1,000. In Dublin and suburbs 138 deaths were registered, including diphtheria 1, enteric fever 1, measles 2, scarlet fever 1, and 35 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended June 24, 1905, correspond to an annual rate of 15.1 per 1,000 of the population, which is estimated at 1,749,917. The lowest rate of mortality was recorded in Leith, viz, 9.6, and the highest in Greenock, viz, 18.6, per 1,000. The aggregate number of deaths registered from all causes was 508, including diphtheria 4, measles 18, scarlet fever 2, and 18 from whooping cough.

SPAIN—Barcelona.—Ten days ended June 20, 1905. Estimated population, 600,000. Total number of deaths, 403, including diphtheria 4, enteric fever 3, measles 12, whooping cough 8, smallpox 5, and 19 from tuberculosis.

Cholera, yellow fever, plague, and smallpox, June 30 to July 21, 1905.

[Reports received by the Surgeon-General, Public Health and Marine-Hospital Service, from American consuls through the Department of State, and from other sources.]

[For reports received from December 30, 1904, to June 30, 1905, see PUBLIC HEALTH REPORTS for June 30, 1905.]

[NOTE.—In accordance with custom, the tables of epidemic diseases are terminated semiannually and new tables begun.]

CHOLERA.

Place.	Date.	Cases.	Deaths.	Remarks.
India:				
Calcutta.....	May 27-June 10..	14	
Madras.....	June 3-9.....	1	

YELLOW FEVER.

Place.	Date.	Cases.	Deaths.	Remarks.
Africa:				
Goree-Dakar	May 31	1	
Brazil:				
Rio de Janeiro	May 27-June 11..	85	34	
British Honduras:				
Belize	June 30-July 6...	1	1	
Ecuador:				
Guayaquil.....	June 6-20.....	7	
Guatemala:				
Livingston	June 10-July 6 ..	12	5	
Honduras:				
Puerto Cortez	June 15-July 6...	24	8	
San Pedro	July 6	Epidemic.
Mexico:				
Tehuantepec	June 24-July 1...	1	1	
Veracruz (Coatzacoalcos, Tierra Blanca, and Veracruz).	June 18-July 11..	6	5	
Panama:				
Colon.....	June 16-July 1...	12	4	
Corozal	June 16-22	2	
Empire	June 16-22	1	
La Boca	June 16-22	2	
Panama	June 16-28	16	4	
Paraiso	June 23-28	1	
Venezuela:				
Maracaibo.....	June 22.....	Present